

Remarks on Robin's and Nicolas Inequalities

Boris A. Kupershmidt
The University of Tennessee Space Institute,
Tullahoma, TN 37388
bkupershmidt@utsi.edu

April 20, 2009

Abstract

Nicolas Conjecture is disproved. The Robin Conjecture follows.

To V. I. Arnold, the greatest mathematician of all times.

I believe this to be false. There is no evidence whatever for it (unless one counts that it is always nice when any function has only real roots). One should not believe things for which there is no evidence. In the spirit of this anthology I should also record my feelings that there is no imaginable reason why it should be true.

J. E. Littlewood on Riemann's Conjecture

... We are all in our own eyes a failure: after all, we haven't proved Fermat's Last Theorem, nor Riemann's Conjecture.

Mary L. Cartwright

1 Introduction

The Nicolas Conjecture [Nic 1983] states that

$$\frac{\mathcal{N}_k}{\varphi(\mathcal{N}_k)} > e^\gamma \log \log \mathcal{N}_k, \quad k \geq 1, \quad (0)$$

where:

$$\mathcal{N}_k = \prod_{i=1}^k p_i,$$

p_i is the prime number $\#i$, φ is the Euler phi-function, and $\gamma = 0.57\dots$ is the Euler constant. For more details, see the beautiful paper [CLM 2006], where it was proven that

$$\prod_{i=1}^k (p_i + 1)/p_i < e^\gamma \log \log \mathcal{N}_k^2, \quad k > 4,$$

in contrast to (1):

$$\prod_{i=1}^k p_i/(p_i - 1) > e^\gamma \log \log \mathcal{N}_k, \quad k \geq 1.$$

Crucially, Nicolas proved that if his Conjecture is *not* true then the inequality (1) is both true and untrue infinitely often. Thus, it's enough to establish it for $k \gg 1$, i.e. for k large enough.

The Robin's inequality, equivalent to Nicolas one, is:

$$\frac{\sigma(n)}{n} < e^\gamma \log \log n, \quad n \geq 5041, \quad (1)$$

where $\sigma(n) = \sigma_1(n) = \sum_{d|n} d$ is the sum of divisors of n . [Rob 1984].

As is clear from the table of exceptions ≤ 5040 in [CLM 2006], (1) can be replaced by:

$$\frac{\sigma(n)}{n} < e^\gamma \log \log n, \quad \omega(n) > 4, \quad (2)$$

where $\omega(n)$ is the number of distinct prime divisors of n .

CLMS also mention that it's enough to consider only those n which are in Hardy-Ramanujan form: if $\omega(n) = m$,

$$n = p_1^{e_1} \dots p_m^{e_m}, \quad (3)$$

then

$$e_1 \geq e_2 \geq \dots \geq e_m.$$

Still, the inequality (1) has arbitrary many parameters: the e_i 's. This is not conducive to a proof. The Nicolas conjecture offers better chances.

2 The Method

Thus, in handling the Nicolas inequality, we need

$$\log LHS \stackrel{?}{>} \log RHS,$$

where

$$\log LHS = \sum_{i=1}^m \log \left(1 + \frac{1}{p_i - 1} \right), \quad (4)$$

$$\log RHS = \gamma + \log \log \theta(p_m). \quad (5)$$

Now,

$$\begin{aligned}\theta(p_m) &\leq p_m \left(1 + \frac{\eta_s}{\log^s p_m}\right), \quad s = 1, 2, 3 \Rightarrow \\ \log \theta(p_m) &\leq \log p_m + \frac{\eta_s}{\log^s p_m} = \log p_m \left(1 + \frac{\eta_s}{\log^{s+1} p_m}\right) \Rightarrow \\ \log \log \theta(p_m) &\leq \log \log p_m + \frac{\eta_s}{\log^{s+1} p_m},\end{aligned}\tag{6}$$

and we are going to look, beyond first 2 terms, $\gamma + \log \log p_m$, at the series in $\frac{1}{\log p_m}$. Thus,

$$\log RHS = \log \log p_m + \gamma + \frac{\eta_3}{\log^4 p_m} + \dots\tag{7}$$

For the LHS, we have:

$$\begin{aligned}\sum_{i=1}^n \log \left(1 + \frac{1}{p_i - 1}\right) &= \sum_{i=1}^m \left\{ \left[\log \left(1 + \frac{1}{p_i - 1}\right) - \frac{1}{p_i} \right] + \frac{1}{p_i} \right\} = \\ &= \sum_{i=1}^m \left[\log \left(1 + \frac{1}{p_i - 1}\right) - \frac{1}{p_i} \right] + \sum_{i=1}^m \frac{1}{p_i} = \sum_{i=1}^m \frac{1}{p_i} + \sum_{i=1}^{\infty} \left[\log \left(1 + \frac{1}{p_i - 1}\right) - \frac{1}{p_i} \right] - \\ &\quad - \sum_{m+1}^{\infty} \left[\log \left(1 + \frac{1}{p_i - 1}\right) - \frac{1}{p_i} \right] \left(\left[\text{Fin 2003} \right], \text{p.95} \right) = \\ &= \sum_{i=1}^m \frac{1}{p_i} + (\gamma - M) - \sum_{m+1}^{\infty} \left[\log \left(1 + \frac{1}{p_i - 1}\right) - \frac{1}{p_i} \right],\end{aligned}\tag{8}$$

where ([Fin 2003], p. 95)

$$\sum_{i=1}^m \frac{1}{p_i} = \log \log p_m + M + o(1),\tag{9}$$

$$M = 0.261497\dots\tag{10}$$

Let's dispose now of \sum_{m+1}^{∞} - term. We have:

$$\log \left(1 + \frac{1}{p-1}\right) - \frac{1}{p} \sim \frac{1}{p-1} - \frac{1}{p} \sim \frac{1}{i^2 \log^2 i}$$

and

$$\int_{m+1}^{\infty} \frac{dx}{x^2 \log^2 x} \sim -\frac{1}{x \log^2 x} \Big|_{m+1}^{\infty} \sim \frac{1}{m \log^2 m},$$

and we are going to count only m -free terms in comparing the log LHS with the log RHS.

Thus, the first two leading terms: $\log\log p_m$ and γ - being equal on both sides, we are comparing

$$\sum_{i=1}^m \frac{1}{p_i} - \log\log p_m - M \text{ and } 0, \quad (11)$$

at least modulo

$$\frac{1}{\log^3 m}. \quad (12)$$

We need the first nonzero term in the log LHS, in (11), being > 0 (or < 0 , as the case may be).

3 The Proof

We calculate modulo $1/m^2$ and modulo $1/m\log^4 m$. Our coefficients are polynomial functions in $w = \log\log m$.

Lemma 13.

$$\log(m+1) = \log m + \frac{1}{m}. \quad (14)$$

Proof.

$$\log(m+1) = \log\left[m\left(1 + \frac{1}{m}\right)\right] = \log m + \log\left(1 + \frac{1}{m}\right) \equiv \log m + \frac{1}{m} \quad \blacksquare$$

Lemma 15.

$$\log\log(m+1) = \log\log m + \frac{1}{m\log m}. \quad (16)$$

Proof.

$$\begin{aligned} \log\log(m+1) &= \log\left(\log m + \frac{1}{m}\right) \equiv \log\left[\log m\left(1 + \frac{1}{m\log m}\right)\right] = \\ &= \log\log m + \frac{1}{m\log m}. \quad \blacksquare \end{aligned} \quad (1)$$

Let $C = C(w)$, $w = \log\log m$. Call it C_m .

Lemma 17.

$$\frac{C_m}{\log^k m} - \frac{C_{m+1}}{\log^k(m+1)} = \frac{kC - C'}{m\log^{k+1} m}. \quad (18)$$

Proof. We have:

$$\begin{aligned} LHS &= \frac{C}{\log^k m} - \frac{C + C' \frac{1}{m \log m}}{\left(\log m + \frac{1}{m}\right)^k} = \frac{1}{\log^k m \left(\log^k m + \frac{k}{m} \log^{k-1} m\right)} \text{ times :} \\ & C \left(\log^k m + \frac{k}{m} \log^{k-1} m\right) - \left(C + C' \frac{1}{m \log m}\right) \log^k m = \\ & = \frac{\log^{k-1} m}{m} (kC - C'). \end{aligned}$$

Altogether,

$$LHS = \frac{1}{m \log^{k+1} m} (kC - C'). \quad \blacksquare$$

Lemma 19.

$$\frac{1}{p_{m+1}} = \frac{1}{p_m}. \quad (20)$$

Proof. Set

$$p_m = m f(m), \quad (21)$$

where

$$\begin{aligned} f(m) &= \log m + (\log \log m - 1) + \frac{\log \log m - 2}{\log n} + \dots = \\ &= \log m + \sum_{i \geq 0} \frac{P_i}{\log^i m}. \end{aligned} \quad (22)$$

Denote $P(m+1) = \tilde{P}$. Then

$$\frac{1}{p_m} - \frac{1}{p_{m+1}} = \frac{1}{p_m p_{m+1}} (p_{m+1} - p_m) \sim \frac{1}{m^2 \log^2 m} (p_{m+1} - p_m), \quad (23)$$

and

$$\begin{aligned} p_{m+1} - p_m &= (m+1)f(m+1) - mf(m) = \\ &= m[f(m+1) - f(m)] + f(m+1) \equiv m[f(m+1) - f(m)], \end{aligned}$$

and

$$f(m+1) - f(m) = \log(m+1) - \log m + \sum_{i \geq 1} \left(\frac{\tilde{P}_i}{\log^i(m+1)} - \frac{P_i}{\log^i m} \right).$$

But

$$\log(m+1) - \log m = \frac{1}{m},$$

and by (18),

$$\frac{\tilde{P}_i}{\log^i(m+1)} - \frac{P_i}{\log^i m} = \frac{P'_i - iP_i}{m \log^{i+1} m}.$$

Together with (23), this proves (20). ■

We are interested in $C = C_1, D = C_2$, such that

$$\sum_{i=1}^m \frac{1}{p_i} = \log \log p_m + M + \frac{C}{\log m} + \frac{D}{\log^2 m} + \dots$$

We use induction on m to find C and D . We have:

$$\begin{aligned} \sum_{i=1}^m \frac{1}{p_i} + \frac{1}{p_{m+1}} &= \frac{1}{p_{m+1}} + \log \log p_m + M + \frac{C}{\log m} + \frac{D}{\log^2 m} = \\ &= \log \log p_{m+1} + M + \frac{\tilde{C}}{\log(m+1)} + \frac{\tilde{D}}{\log^2(m+1)}, \end{aligned} \quad (24)$$

or

$$\left(\frac{C}{\log m} - \frac{\tilde{C}}{\log(m+1)} \right) + \left(\frac{D}{\log^2 m} - \frac{\tilde{D}}{\log^2(m+1)} \right) = (\log \log p_{m+1} - \log \log p_m) - \frac{1}{p_m}. \quad (25)$$

By (18), this is:

$$\frac{C - C'}{m \log^2 m} + \frac{2D - D'}{m \log^3 m} = (\log \log p_{m+1} - \log \log p_m) - \frac{1}{p_m}. \quad (26)$$

Let's start with the $1/p_m -$ term. We have:

$$\begin{aligned} -\frac{1}{p_m} &= -\frac{1}{m \log m} \left(\frac{1}{1 + \frac{P_0}{\log m} + \frac{P_1}{\log^2 m}} \right) = \\ &= -\frac{1}{m \log m} \left(1 - \frac{P_0}{\log m} - \frac{P_1}{\log^2 m} + \frac{P_0^2}{\log^2 m} \right). \end{aligned} \quad (27)$$

Lemma 28.

$$\log p_{m+1} - \log p_m \sim \frac{1}{m} \left(1 + \frac{1}{\log m} \right) - \frac{P'_0 - P_0}{m \log^2 m}. \quad (29)$$

Proof. We have:

$$\begin{aligned}
LHS &= \log \frac{p_{m+1}}{p_m} = \log \left[\frac{m+1}{m} \frac{f(m+1)}{f(m)} \right] = \frac{1}{m} + \log \frac{f(m+1)}{f(m)} = \\
&= \frac{1}{m} + \log \left\{ \frac{\log(m+1)}{\log m} \left[\frac{1 + \frac{\tilde{P}_0}{\log(m+1)} + \frac{\tilde{P}_1}{\log^2(m+1)} + \dots \right] \right\} = \\
&= \frac{1}{m} + \frac{1}{m \log m} + \log \left[\begin{array}{c} \dots \\ \dots \end{array} \right] = \\
&= \frac{1}{m} \left(1 + \frac{1}{\log m} \right) + \log \left[1 + \frac{\tilde{P}_0}{\log(m+1)} + \frac{\tilde{P}_1}{\log^2(m+1)} + \dots \right] - \log \left[1 + \frac{P_0}{\log m} + \frac{P_1}{\log^2 m} + \dots \right] = \\
&= \frac{1}{m} \left(1 + \frac{1}{\log m} \right) + \frac{\tilde{P}_0}{\log(m+1)} - \frac{P_0}{\log m} \quad [\text{by (18)}] = \\
&= \frac{1}{m} \left(1 + \frac{1}{\log m} \right) - \frac{P'_0 - P}{m \log^2 m}. \quad \blacksquare
\end{aligned}$$

Lemma 30.

$$\frac{1}{\log p_m} = \frac{1}{\log m} \left(1 - \frac{\log \log m}{\log m} \right) + O\left(\frac{1}{\log^3 m} \right). \quad (31)$$

Proof. We have:

$$\begin{aligned}
\frac{1}{\log p_m} &= \frac{1}{\log m + \log f(m)} = \frac{1}{\log m} \frac{1}{\left[1 + \frac{1}{\log m} \log f(m) \right]} = \\
&= \frac{1}{\log m} \left(1 - \frac{\log \log m}{\log m} \right) \quad \blacksquare
\end{aligned}$$

Lemma 32.

$$\log \log p_{m+1} - \log \log p_m = \frac{1}{m \log m} \left(1 + \frac{1}{\log m} \right) + \frac{P'_0 - P_0}{m \log^3 m} - \frac{\log \log m}{m \log^2 m} \left(1 + \frac{1}{\log m} \right). \quad (33)$$

Proof. We have:

$$\begin{aligned}
LHS &= \log \frac{\log p_{m+1}}{\log p_m} = \log \left(1 + \frac{\log p_{m+1} - \log p_m}{\log p_m} \right) = \\
&= \frac{\log p_{m+1} - \log p_m}{\log p_m} \quad [\text{by (29), (31)}] = \\
&= \frac{1}{\log m} \left(1 - \frac{\log \log m}{\log m} \right) \left\{ \frac{1}{m} \left(1 + \frac{1}{\log m} \right) - \frac{P'_0 - P_0}{m \log^2 m} \right\} = \\
&= \frac{1}{m \log m} \left(1 + \frac{1}{\log m} \right) - \frac{P'_0 - P_0}{m \log^3 m} - \frac{\log \log m}{m \log^2 m} \left(1 + \frac{1}{\log m} \right). \quad \blacksquare
\end{aligned}$$

Collecting all Lemmas, we rewrite (25) as

$$\begin{aligned}
& \frac{C - C'}{m \log^2 m} + \frac{2D - D'}{m \log^3 m} = \\
& = \frac{1}{m \log m} \left(1 - \frac{P_0}{\log m} + \frac{P_0^2 - P_1}{\log^2 m} \right) + \\
& + \frac{1}{m \log m} \left(1 + \frac{1}{\log m} \right) - \frac{P'_0 - P_0}{m \log^3 m} - \frac{\log \log m}{m \log^2 m} \left(1 + \frac{1}{\log m} \right). \quad (34)
\end{aligned}$$

Thus,

$$\begin{aligned}
& \frac{1}{m \log m} - \text{coefficient is } 0; \\
& \frac{1}{m \log^2 m} - \text{coefficient is :} \\
& P_0 + 1 - \log \log m = 0 \text{ because } P_0 = \log \log m - 1; \\
& \frac{1}{\log^3 m} - \text{coefficient is :} \\
& -P_0^2 + P_1 - P'_0 + P_0 - \log \log m < 0 \text{ for } m \gg 0. \text{ Thus,} \\
& \quad \quad \quad LHS < RHS. \quad (35)
\end{aligned}$$

The Nicolas Conjecture is thus disproved, but the strengthened Robin Inequality, is established:

$$\prod_{i=1}^m \frac{p_i}{\varphi(p_i)} < e^\gamma \log \log \mathcal{N}_m.$$

Notice that Nicolas Inequality is disproved in a way that contradicts the Nicolas result, that if his inequality is wrong, then it's wrong *and right* infinitely often. Note that the equation

$$kF - F' = p(w),$$

where p is a polynomial, has trivially a unique polynomial solution in $w = \log \log m$.

4 Acknowledgment 1

Although not apparent, the paper is a result of hundreds of computer experiments, and it has an invisible hero: the PARI-GP program, a free to the mathematical community wonderful tool for computer experiments. My deep gratitude to the developers of PARI.

Nicholas Lynn was instrumental in discovering widespread order among the primes, by running many numerical experiments for me on Mathematica. Later Tony Saad had replaced him. Joshua Kuperchmidt set me up with PARI, and helped both with its quirks and the actual programming.

I thank Remi Engels for technical assistance with Robin's article, and Keith Briggs for sending me his inspiring paper.

Last but not least, over the years of struggles with the problem, with me often concluding that it's unsolvable, K. C. Reddy held unshakably to the firm believe that I'll conquer it. Why he was so certain, is a mystery to me.

5 Acknowledgment 2

It would be churlish of me not to mention the wonderfully well written and remarkably exciting and stimulating paper [CLMS 2006].

6 Acknowledgment 3

I'm grateful to Pieter Moore for the careful reading of the first version of this paper and constructing a counterexample.

I'm also deeply indebted to Patrick Solé for his encouragement after reading the fatally flawed 1st version of the Manuscript.

7 Acknowledgment 4

The dim-witted editors of the Archive had removed the 1st version of the Manuscript "for inappropriate language". When I appealed their thuggish action to Grinspurg, Kuperberg, and Tao, I got not response.

8 Acknowledgment 5

This paper might have never been finished were it not for the vaguely unthreatening letter from Thi-Eve Jao, the Editor-in-Chief of the International Journal of Chinese Mathematics, in which he politely explained that idleness of people like me threatens the livelihood of people like him, and implored me to finish the proof of the Nicolas conjecture as expeditiously as possible, but no later than the opening of the next Congress of the Chinese Communist Gangsters Party. There was nothing else to do but to comply with so exquisitely framed request.

References

- [Bri 2006] Briggs, K., *Abundant Numbers and the Riemann Hypothesis*, J. Exper. Math. **15** (2006) 251-256.
- [CLM 2006] Choie, Y.-J., Lichiardopol, N., Moree, P., and Solé, *On Robin's criterion for the Riemann Hypothesis*, arXiv:math.NT/0604314; J. Théor. Nombres Bordeaux 19 (2007), no. 2, 357–372.

- [Dus 1998] Dusart, P., *Autour de la fonction qui compte le nombre de nombres premiers*, Theses, Universite de Limoges, France (1998) http://www.unilim.fr/laco/theses/1998/T1998_01.pdf
- [Fin 2003] Finch, S. R., *Mathematical constants*, Encyclopedia of Mathematics and its Applications **94**, (Cambridge University Press, Cambridge, 2003).
- [Lan 1906] Landau, E., *Handbuch der Lehre von der Verteilung der Primzahlen*, Leipzig (1909).
- [Nic 1983] Nicolas, J. - L., *Petite valeurs de la fonction d'Euler*, J. Number Theory **17** (1983) 375-388.
- [Nic 1988] Nicolas, J. L., *Bornes Effectives Pour Certaines Fonctions Arithmetiques*, Colloque de Théorie Analytique des Nombres "Jean Coquet" (Marseille, 1985) 91-99, Publ. Math. Orsay, 88-02, Univ. Paris XI, Orsay, 1988.
- [PAR 2006] PARI/GP, Version 2.3.0, Bordeaux (2006), available from <http://pari.math.u-bordeaux.fr>
- [Rob 1984] Robin, G., *Grandes Valeurs de la fonction somme des diviseurs et hypothese de Riemann*, J. Math. Pures Appl. (9) **63** (1984) 187-213.
- [RSh 1962] Rosser, J. B., and Shoenfeld, L., *Approximate Formulas for Some Functions of Prime Numbers*, Illinois J. Math. **6** (1962) 69-94.
- [RSh 1975] Rosser, J. B., and Shoenfeld, L., *Sharper Bounds for the Chebyshev Functions $\theta(x)$ and $\psi(x)$* , Math. Comp. **29** (1975) 243-269.
- [YYa 2009] Yaweeccan, X. M., and Yahodoo, Y.Z., *The Only True and Definitive Proof Of The Great Chinese Theorem Formerly Known As Riemann's Hypothesis*, Asian J. of Pilfered Math. (to be announced as soon as possible, but no later than the last year.)